


REPORT DOCUMENTATION PAGE			Form Approved OMS No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget Paperwork Reduction Project (0704-0188), DC 20503.				
1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE 2/18/97		
		3. REPORT TYPE AND DATES COVERED Final 1 December 1993 - 30 November 1996		
4. TITLE AND SUBTITLE Large Amplitude Motions in Polyatomic Molecule Spectra: Intramolecular Vibrational Redistribution and Isomerization		5. FUNDING NUMBERS 61102 F 2303 B1 AFOSR-TR-96 97 0128		
6. AUTHOR(S) Robert W. Field and Robert J. Silbey				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Massachusetts Institute of Technology 77 Massachusetts Avenue Cambridge, MA 02139-4307		8. PERFORMING ORGANIZATION REPORT NUMBER 4		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NC Building 410 Bolling Air Force Base Washington, DC 20332-6448 NL		10. SPONSORING/MONITORING AGENCY REPORT NUMBER F49620-94-1-0068		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION IS UNLIMITED		12b. DISTRIBUTION CODE 19970602 058		
13. ABSTRACT (Maximum 200 words) Through Stimulated Emission Pumping (SEP) studies of highly excited vibrational levels of the electronic ground state of HCP, the spectroscopic signatures of bond-breaking isomerization ($\text{HCP} \leftrightarrow \text{HPC}^+$) were detected and interpreted for the first time. Two novel spectroscopic schemes, UV excited-IR detected fluorescence excitation spectroscopy and UV excited-Auger detected Laser Excited Metastable (LEM) spectroscopy, prove the existence of a Gateway Mediated Inter-System Crossing (GMISC) mechanism in acetylene. In GMISC, one special vibronic state couples the single bright state to a dense manifold of dark states. The complementarity of LEM and Laser Induced Fluorescence (LIF) spectroscopies was found to be ideally suited for study of ISC, because LEM is blind to short-lived radiating states and LIF is blind to metastable states. A pattern recognition technique, Extended Cross Correlation (XCC), which is capable of picking out multiple patterns that are repeated in multiple spectra without any prior knowledge of the forms of the patterns, was applied with extraordinary success to several types of spectra. Among these was a data set consisting of electron beam excited infrared spectra of CO, recorded by Dr. Stephen Lipson and colleagues at the Air Force Phillips Laboratory. All of the severely overlapped features were "unzipped," assigned, and independently represented by time dependent populations. The central theme in each of these projects is the development of experimental methods, theoretical models, and pattern-recognition schemes capable of extracting previously inaccessible information from spectra of highly excited molecules.				
14. SUBJECT TERMS Laser Spectroscopy, Potential Energy Surface Barriers, Intramolecular Vibrational Redistribution, Isomerization, Triplet States, Intersystem Crossing		15. NUMBER OF PAGES 16. PRICE CODE		
17. SECURITY CLASSIFICATION UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL (UNLIMITED) SAR (SAME AS REPORT)	

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE
COPY FURNISHED TO DTIC
CONTAINED A SIGNIFICANT
NUMBER OF PAGES WHICH DO
NOT REPRODUCE LEGIBLY.**

Final Report

to

Air Force Office of Scientific Research

**Large Amplitude Motions in Polyatomic Molecule
Spectra: Intramolecular Vibrational Redistribution
and Isomerization
F49620-94-1-0068**

Presented by:

Robert W. Field
Professor of Chemistry
and
Robert J. Silbey
Professor of Chemistry

Massachusetts Institute of Technology
77 Massachusetts Avenue
Cambridge, Massachusetts 02139-4307

Period: 1 December 1993 - 30 November 1996

TABLE OF CONTENTS

A.	Final Report	1
1.	LIST OF PERSONNEL	3
a.	Graduate students	3
b.	Postdocs and Visitors.....	3
c.	Undergraduate Students.....	4
d.	Collaborators.....	4
2.	Report on RESEARCH PROJECTS.....	5
a.	HCP Isomerization.....	5
b.	Triplet States of Acetylene.....	6
c.	Pattern Recognition.....	6
d.	Resonance Enhanced Multi-Photon Ionization Apparatus	8
e.	Ionization Energies of CaF and BaF.....	8
B.	Publications (since 1986) Resulting from AFOSR Support	9
C.	Ph.D. Theses (since 1986) Resulting from AFOSR Support.....	13

A. Final Report 1 December 1993 - 30 November 1996

1. LIST of Personnel

a. Graduate Students

Jonathan Bloch	(Ph.D. 5/96) ¹
Jason Clevenger ¹	
Kevin Cunningham	
Ilia Dubinsky ³	
Christopher M. Gittins	(Ph.D. 2/95) ¹
Matthew Jacobson ⁴	
Neme Nnolim	(M.S. 9/95)
Jonathan P. O'Brien ²	
Bhavani Rajaram	(Ph.D. 1/95)
Leah Ruslen ²	
Gunther Schmid ⁶	
Michelle Silva	
Stephani Ann B. Solina	(Ph.D. 5/96) ^{2,5}
Jianghong Wang	(M.S. 5/94)

1. Primary support NSF
2. Primary support DOE
3. Primary support J. Steinfeld NASA
4. DOD Predoctoral Fellow
5. NSF Predoctoral Fellow
6. Primary Support R. Silbey NSF

b. Postdocs and Visitors

Dr. Stephen Coy ¹	
Dr. Stephen Drucker	
Dr. Brian Gilbert	(now at Coastal Carolina Univ.) ⁴
Dr. Haruki Ishikawa	(now at Tohoku University) ²
Dr. Bing Ji	(now at Air Products Corp.) ⁵
Prof. Marsha Lester	(U. of Pennsylvania) ³
Dr. David Moss	(Boston University) ⁶
Prof. William Polik	(Hope College) ³
Prof. Richard Redington	(Texas Tech University)
Prof. Curt Wittig	(U. of Southern California) ³

1. Consultant
2. JSPS Fellow
3. Sabbatical Visitor
4. Primary support RWF-JIS NASA
5. Primary Support NSF
6. Primary support ACS-PRF

c. Undergraduate Students

Heather Drake
 Ana Isasi
 Jonathan Katz
 Maya Kaushal
 Donald Lucas
 Paresh Patel
 Jennifer Sokol

d. Collaborators

Dr. Michael Davis	Argonne National Lab	pattern recognition
Prof. Edward E. Eyler	U. Connecticut	Pulsed FM
Prof. Robert Gordon	U. Illinois, Chicago	Deperturbation HCl
Prof. Fritz Grein	U. New Brunswick	HCP <i>ab initio</i>
Dr. John Hall	JILA	$^{13}\text{C}_2\text{H}_2$ Isomerization by NICEOHMS
Prof. Joshua Halpern	Howard U.	Cyanogen
Prof. Michel Herman	Bruxelles	HCCH
Prof. Michael Kellman	U. Oregon	pattern recognition
Prof. Kevin Lehmann	Princeton	Intracavity MRS
		Unzipping Isotopic NH_3
Dr. Stephen Lipson	AF Phillips Lab	pattern recognition
Dr. Edmond Murad	AF Phillips Lab	ionization energies
Dr. Eric Rohlfing	DOE, Combustion Research Facility	Unzipping Vinox
Dr. Reinhard Schinke	MPI Gottingen	HCP
Dr. Trevor Sears	Brookhaven National Lab	FM, HCB
Prof. Alec Wodtke	UC Santa Barbara	triplets
Prof. Robert Wyatt	U. Texas	patterns in <i>ab initio</i> spectra

2. REPORT ON RESEARCH PROJECTS

a. HCP Isomerization

This is one of three most important results from this grant.

HCP is the first example in which a bond-breaking isomerization process is sampled by an eigenstate-resolved spectrum. The spectroscopic signature of isomerization, a change in the $\omega_1 : \omega_2 : \dots \omega_{3N-6}$ vibrational resonance structure as manifest in sudden modifications of spectroscopic patterns, is observed in the Stimulated Emission Pumping (SEP) spectrum of the HCP $\tilde{X}^1\Sigma^+$ electronic state (recorded via the \tilde{A}^1A'' and \tilde{C}^1A' intermediate states). The SEP spectra of the HCP \leftrightarrow HPC isomerization process provide examples of several general methods for recognition, analysis, and interpretation of unimolecular isomerization. These include: (i) a sudden onset of (a new class of) spectroscopic perturbations; (ii) a sudden change (departure from normal v_J -scaling) of vibration-rotation fine structure constants; and (iii) the appearance of new classes of vibrational states with anomalously large or small rotational constants. It is essential to be able to follow a long vibrational progression (not necessarily in the normal mode that couples most strongly to the minimum energy isomerization path) continuously up to vibrational energies above the *ab initio* predicted isomerization barrier (or saddle point). It is also essential to have a larger scale spectroscopic pattern, such as the cluster of mutually interacting states that belong to the same value of polyad quantum number(s). In HCP, the 2:1 resonance between the CP stretch (mode 3) and the bend (mode 2), $\omega_3 \approx 2\omega_2$, provided the polyad (polyad quantum number $P = v_2 + 2v_3$) pattern that enabled us to make definitive vibrational assignments of the complicated appearing spectrum at $E_{VIB} \leq 25,000 \text{ cm}^{-1}$.

In a continuing collaboration with Dr. Reinhard Schinke's group at the Max Planck Institut für Stromungsforschung in Gottingen, we first were able to show that all of the dramatic changes observed in the SEP spectrum of HCP were due to the decrease in the H-X (X is the center of mass of CP) stretch from $\sim 5\omega_{\text{bend}}$ to slightly less than $3\omega_{\text{bend}}$ and then back to slightly more than $3\omega_{\text{bend}}$ at linear HPC. This change in resonance structure accounts for the sudden turning on near $v_{\text{bend}} = 32$ of anharmonic ($4\omega_{\text{bend}} \approx \omega_{\text{HX}}$) and Coriolis ($3\omega_{\text{bend}} \approx \omega_{\text{HX}}$) perturbations, the rapid change of all fine structure parameters (defined by second order perturbation theory), and the appearance of an anomalously large-B polyad of $\ell = 0$ substates. Schinke also predicted a class of delocalized states (the "saddle node" SN states), which were subsequently observed by Ishikawa et al in his SEP experiments at Tohoku University. These SN states are the first experimentally observed example of over-the-barrier delocalized states.

The experimental study of HCP at MIT is terminated, but is continuing in Professor Ishikawa's group at Tohoku University. Our collaboration with Ishikawa (experiment) and Schinke (Theory) is continuing. We expect to summarize the new experimental measurements (on the SN states and on the postulated small-B $\ell = 1$ states that are responsible, via Coriolis perturbations, for the large-B $\ell = 0$ states) in several publications, culminating in a review article in *Ann. Revs. Phys. Chem.*

b. Triplet States of Acetylene

This is one of the three most important results from this grant and is the central focus of the continuing grant (F49620-97-1-0040).

We have observed intersystem crossing (ISC) in acetylene by qualitatively new types of experimental schemes: IR detected ($T_{2,3} \rightarrow T_1$)-UV excited ($S_1 \leftarrow S_0$) fluorescence excitation and Auger-detected (T_3), UV-excited "Laser Excited Metastable" (LEM) spectroscopy. A report of the former experiment will appear as a Communication in JCP and a report of the latter experiment has recently been submitted as a Communication to JCP. The unifying theme of both experiments is that it is now possible to characterize the ISC mechanism more completely. Instead of a simple model of one "bright state" coupling to a dense manifold of "dark states," we have proposed a "gateway mediated" ISC mechanism whereby one bright state couples to a single gateway state which in turn couples to a dense manifold of dark states. In the case of S_1 acetylene, the gateway is a single low vibrational level of the T_3 triplet surface (the upper sheet of the Renner-Teller split electronic state that correlates with $\pi_u^3\pi_g^*{}^3\Delta_u$ in linear geometry).

The LEM experiments were carried out in Alec Wodtke's laboratory at UCSB. A supersonic beam of C_2H_2 in H_2 , He, or Ne was excited via various $\tilde{A} - \tilde{X}$ bands, including the $V_0^4K_0^1$ band which terminates in an upper state 200 cm^{-1} above the $C_2H_2 \rightarrow H + CCH(\tilde{X})$ dissociation limit. Metastable species are detected by ejection of an Auger electron from a Au electrode (work function 5.1eV). The high work function of the Au electrode requires T_3 or S_1 character for Auger detection, and the $\sim 200\mu\text{s}$ flight time from excitation to detection restricts the fractional S_1 character to $<1\%$. Therefore LIF and LEM schemes are complementary; one is blind to triplets and the other is blind to singlets. The LEM scheme is able to distinguish intact C_2H_2 metastables from metastable photofragments (C_2H , C_2 , and CH). The observation of intact metastable 200 cm^{-1} above the dissociation limit has startling photochemical consequences. The LEM spectrum exhibits two interference features, which are the universal signature of the gateway mediated mechanism, and in the present case serve to locate the J-value of the level crossing between the bright ($S_1\ 3v_3$) and gateway (T_3) states.

We expect to record a series of higher resolution ($\sim 0.01\text{ cm}^{-1}$) LEM spectra at UCSB in order to better characterize the gateway state. We have nearly completed construction of a complementary electron bombardment supersonic jet source of C_2H_2 metastables at MIT.

c. Pattern Recognition

Modern spectroscopic techniques are capable of generating enormous quantities of high quality (spectral and temporal resolution, quantitative intensities over large dynamic range) spectra. We have developed a special kind of pattern recognition scheme which is capable of picking out patterns that appear in multiple spectra without making any assumptions about the form of the pattern. This scheme, which we call Extended Cross Correlation (XCC) and Extended Auto-Correlation (XAC), has been applied to several classes of spectra: (i) dispersed fluorescence (DF) spectra of $C_2H_2\ \tilde{A} \rightarrow \tilde{X}$; (ii) IR spectra of CO pulse-excited by a kV electron beam; and (iii) FTIR absorption spectra of $NH_3/NDH_2/ND_2H/ND_3$ (recorded at Kitt Peak Solar Observatory) at a series of H : D isotope ratios.

The DF spectra of C_2H_2 are the primary subject of a DOE grant. The SCC method (developed jointly by Steve Coy and Matt Jacobson, supported respectively by AFOSR and a DOD fellowship) has enabled us to "unzip" the spectrum into overlapping polyads, from which we have been able to deduce the nature and rates of the dominant energy flow pathways, which are collectively known as Intramolecular Vibrational Redistribution (IVR). Guided by our AFOSR supported study of the signature of isomerization in HCP, we are about to attempt to unzip the DF spectra of HCCH in the energy region of the acetylene \leftrightarrow vinylidene isomerization barrier. We expect to detect the onset of isomerization via the change in $\omega_1 : \omega_2 : \dots : \omega_{3N-6}$ resonance structure, as revealed by a departure of the polyad structure from vibrational scaling predictions.

A collaboration between Matt Jacobson and Dr. Stephen Lipson at the Air Force Phillips Laboratory is the most important "transition" of this AFOSR project. The following abstract has been submitted for clearance at Phillips Laboratory, and will be the basis for talks by Dr. Lipson (at the Spring meeting of the American Geophysical Union) and Mr. Jacobson (at the 52nd International Symposium on Molecular Spectroscopy, Ohio State University):

Infrared Spectral Data Analysis and Remote Sensing Using Pattern Recognition Algorithms; S.J. Lipson, R.B. Lockwood, D.L. Vitito, W.A.M. Blumberg, P.S. Armstrong, M.P. Jacobson, and R.W. Field.

In remote sensing of atmospheric spectra, the complexity of analyzing data may be compounded by extreme departures from thermodynamic equilibrium, and by spectral overlaps, temporal variation, and lack of knowledge of line-of-sight effects. We have developed algorithms for the analysis of infrared spectral data taken under these conditions, and applied them to the study of highly vibrationally and rotationally excited carbon monoxide observed in atmospheric simulation experiments. The data were taken using the cryogenic-background LABCEDE facility at the Phillips Laboratory. Mixtures of CO, Ar, and N_2 at pressures of 2-40 millitorr were irradiated with a pulsed 4-5 keV electron beam, and the resulting time-resolved infrared CO spectra were obtained with a Michelson interferometer capable of 2 cm^{-1} resolution. The spectra were highly self-absorbed in the $v=1-0$ band, but not in the highly vibrationally and rotationally excited bands. The complex, overlapped, and partially self-absorbed spectra were analyzed using pattern recognition algorithms, and excellent agreement between data and fit was obtained. Vibrational basis sets (patterns) were obtained both from synthetic spectral models and from the XCC (Extended Cross-Correlation) method. A global fit using this combined model was performed, in which the time dependence of the known basis sets and the XCC patterns was determined. Applications of this method include remote sensing of environments in which the optical properties of the line of sight are poorly known, where spectroscopic basis sets are unavailable, or where spectra change with time or viewing angle, such as in the aurora or other highly structured atmospheric scenes.

The IR spectrum of ammonia is very complex, especially for the nonsymmetric ND_2H and NDH_2 isotopomers. Analysis of such a spectrum is enormously simplified when the raw mixed-isotopomer spectrum is separated by a model- and assignment-free method into spectra of separate isotopomers! By recording spectra

of equilibrated sample at several H/D ratios, the XCC scheme unzips the raw spectra into single isotopomer spectra. This work was done by Steve Coy and Jason Clevenger in collaboration with Dr. Linda Brown and Dr. Jack Margolis (Jet Propulsion Laboratory), and Dr. Michael Dulick (Kitt Peak National Observatory). Similar unzipping schemes are under discussion with Professor David Nesbitt (JILA) for H_2O and H_3O^+ , but are better suited for FTIR than cw tunable laser spectroscopy.

d. Resonance Enhanced Multiphoton Ionization (REMPI) Apparatus.

A pulsed jet molecular beam apparatus with REMPI-TOF capabilities has been constructed by Chris Gittins (Ph.D. 2/95). Although so far this apparatus has been used primarily in NSF supported studies of Rydberg states of CaF, CaCl, and BaF, it was constructed and will be available for time-shared use in REMPI-TOF studies of triplet acetylene.

e. Ionization Energies of CaF and BaF.

Mass spectrometric data, recorded by Edmund Murad and James Gardner at the Air Force Phillips, and low principal quantum number Rydberg state data obtained by Jakubek, Harris, and Gittins at MIT, were combined to yield accurate ionization energies. This represents another transition between a university laboratory and an Air Force laboratory.

B. Publications (Since 1986) Resulting from AFOSR Support

The research of the Co-PI's is supported by AFOSR and DOE grants in the general area of the spectra of vibrationally highly excited polyatomic molecules. The distinction between these two funded projects is discussed in the original research proposals. We list below all of our publications relevant to the research described in the present project; AFOSR support is denoted by (+) and DOE support by (++).

- ++ G.J. Scherer, Y. Chen, R.L. Redington, J.L. Kinsey, and R.W. Field, "An Unsuspected Fermi Perturbation in the Acetylene \tilde{A}^1A_u $3\nu_3$ Level", J. Chem. Phys. 85, 6315-6323 (1986).
- + G. Pichler, J.T. Bahns, K.M. Sando, W.C. Stwalley, D.D. Konowalow, Li Li, R.W. Field, and W. Müller, "Electronic Assignments of the Violet Bands of Sodium", Chem. Phys. Lett. 129, 425-428 (1986).
- ++ J.P. Pique, Y. Chen, R.W. Field and J.L. Kinsey, "Chaos and Dynamics on 30fs-1ns Time Scales in Vibrationally Excited Acetylene: Fourier Transform of Stimulated Emission Pumping Spectrum", Phys. Rev. Letts. 58, 475-478 (1987).
- +,++ C.E. Hamilton, J.L. Kinsey, and R.W. Field, "Stimulated Emission Pumping: New Methods in Spectroscopy and Molecular Dynamics," Ann. Rev. Phys. Chem. 37, 493-524 (1986).
- + F. Temps, S. Halle, P.H. Vaccaro, R.W. Field, and J.L. Kinsey, "Collisional Energy Transfer in Highly Vibrationally Excited H_2CO (\tilde{X}^1A_1)" J. Chem. Phys. 87, 1895-1897 (1987).
- + P.H. Vaccaro, F. Temps, S. Halle, J.L. Kinsey, and R.W. Field, "Polarization-Detected Transient Gain Studies of Relaxation Processes in $\nu_4 = 1$ \tilde{A}^1A_2 Formaldehyde- h_2 ", J. Chem. Phys. 88, 4819-4833 (1988).
- + R.L. Redington, Y. Chen, G.J. Scherer, and R.W. Field, "Laser Fluorescence Excitation Spectrum of Jet-Cooled Tropolone: The $\tilde{A}^1B_2 - \tilde{X}^1A_1$ System", J. Chem. Phys. 88, 627-633 (1988).
- + F. Temps, S. Halle, P.H. Vaccaro, R.W. Field, and J.L. Kinsey, "Vibrationally Excited Formaldehyde: The Relationship between Vibrational Structure and Collisional Properties", Faraday Discussion on Molecular Vibrations, 1987, J. Chem. Soc. Faraday Trans. 2, 84, 1457-1482 (1988).
- ++ J.-P. Pique, Y. Chen, R.W. Field, and J.L. Kinsey, "Laser Spectroscopy and Quantum Chaos: An Example Through the Fourier Transform of a Stimulated Emission Pumping Spectrum of C_2H_2 at Very High Vibrational Energy", Conference Laser M2P, 1987, J. de Physique C7, 655-657 (1987).
- ++ J.P. Pique, Y. Chen, R.W. Field, J.L. Kinsey, and M. Lombardi, "New Order Out of the Chaotic Bath of Highly Excited Vibrational States of C_2H_2 ", Berichte der Bunsengesellschaft für Physikalische Chemie 92, 422-424 (1988).
- +,++ R.W. Field, "High Resolution Spectroscopy of Small Molecules", Conference Laser M2P, 1987, J. de Physique C7, 17-28 (1987).
- ++ Y. Chen, D.M. Jonas, C.E. Hamilton, P.G. Green, J.L. Kinsey, and R.W. Field, "Acetylene: Isomerization and Dissociation," Berichte der Bunsengesellschaft für Physikalische Chemie, 92 329-336 (1988).
- ++ J.P. Pique, Y.M. Engel, R.D. Levine, Y. Chen, R.W. Field, and J.L. Kinsey, "Broad Spectral Features in the Stimulated Emission Pumping Spectrum of Acetylene", J. Chem. Phys. 88, 5972-5974 (1988).
- +,++ M.H. Alexander, P. Andresen, R. Bacis, R. Bersohn, F.J. Comes, P.J. Dagdigian, R.N. Dixon, R.W. Field, G.W. Flynn, K.-H. Gericke, B.J. Howard, J.P. Huber, D.S. King, J.L. Kinsey, K. Kleinermanns, A.C. Luntz, A.J. MacCaffery, B. Pouilly, H. Reisler, S. Rosenwaks, E. Rothe, M. Shapiro, J.P. Simons, R. Vasudev, J.R. Wiesenfeld, C. Wittig, and R.N. Zare, "A Nomenclature for Λ -Doublet Levels in Rotating Linear Molecules", J. Chem. Phys. 89, 1749-1753 (1988).

- + R.L. Redington and R.W. Field, "Laser Fluorescence Excitation Band Profiles of Jet-Cooled Tropolone", *Spectrochim. Acta* 45A 41-46 (1989).
- + P.H. Vaccaro, A. Zabludoff, M. Carrera-Patiño, J.L. Kinsey, and R.W. Field, "High Precision Dipole Moments in A^1A_2 Formaldehyde Determined via Stark Quantum Beat Spectroscopy", *J. Chem. Phys.* 90, 4150-4167 (1989).
- + ,++ R.W. Field, "Pathologically Complex Spectra of Small Molecules: High Resolution Techniques for Displaying Vibrational Chaos and Electronic Simplicity", Conference on Quantum Electronics and Laser Science, 1989 Technical Digest Series, Vol. 12 (Optical Society of America, Washington, DC), pp. 160-162 (Talk THHH2), 1989.
- + ,++ R.W. Field, "Statistical Spectroscopy: Insight or Nonsense?", ILS-IV, 1988, Am. Instit. of Phys. Conf. Proc. 191, 673-682 (1988).
- ++ P.G. Green, J.L. Kinsey, and R.W. Field, "A New Determination of the Dissociation Energy of Acetylene", *J. Chem. Phys.* 91, 5160-5163 (1989).
- + F. Temps, S. Halle, P.H. Vaccaro, R.W. Field, and J.L. Kinsey, "Collisional Relaxation of H_2CO (A^1A_2 , $v_4 = 1$, $J_{K_a, K_c} = 13_{2,12}$) by He, Ar, Xe, and N_2 ", *J. Chem. Phys.* 91, 1008-1011 (1989).
- ++ Y. Chen, D.M. Jonas, J.L. Kinsey, and R.W. Field, "High Resolution Spectroscopic Detection of Acetylene \leftrightarrow Vinylidene Isomerization by Spectral Cross-Correlation", *J. Chem. Phys.* 91, 3976-3987 (1989).
- ++ J. Vander Auwera, T.R. Huet, M. Herman, C. Hamilton, J.L. Kinsey, and R.W. Field, "The \tilde{A}^1A' Electronic State of Monodeuterated Acetylene", *J. Mol. Spectrosc.* 137, 381-395 (1989).
- ++ C.E. Hamilton, R.W. Field, T.R. Huet, and M. Herman, "Low Energy Vibrational Levels in the \tilde{A}^1A_u Electronic State of C_2D_2 ", *J. Mol. Spectrosc.* 137, 427-429 (1989).
- + D.M. Jonas, X. Zhao, K. Yamanouchi, P.G. Green, G.W. Adamson, and R.W. Field, "High Resolution VUV Fluorescence Excitation and Predissociation of \tilde{A}^1A' HCN", *J. Chem. Phys.* 92, 3988-3989 (1990).
- + R.L. Redington, T.E. Redington, M.A. Hunter, and R.W. Field, " $\tilde{A}^1B_2 - \tilde{X}^1A_1$ 26_0^7 Transitions of ^{18}O -Enriched Tropolone", *J. Chem. Phys.* 92, 6456-6462 (1990).
- + Y.-T. Chen, D.M. Watt, R.W. Field, and K.K. Lehmann, "Observation of Highly Vibrationally Excited $\tilde{X}^1\Sigma^+ HCP$ by Stimulated Emission Pumping Spectroscopy", *J. Chem. Phys.* 93, 2149-2151 (1990).
- ++ Y. Chen, S.D. Halle, D.M. Jonas, J.L. Kinsey, and R.W. Field, "Stimulated Emission Pumping Studies of Acetylene $\tilde{X}^1\Sigma_g^+$ in the 11,400-15,700 cm^{-1} Region: the Onset of Mixing", *J. Opt. Soc. Am. B* 7, 1805-1815 (1990).
- ++ C. Kittrell and B.A. Garetz, "Analysis of the $D^1\Delta - X^1\Sigma^+$ Transition in CO Observed by Two-Photon Excitation", *Spectrochim. Acta* 45A, 31-40 (1989).
- + D.M. Jonas, "Spin Statistics: An Error in Landau and Lifschitz' Quantum Mechanics", *J. Chem. Phys.* 90, 5563-5565 (1989).
- + S.L. Coy, R. Hernandez, and K.K. Lehmann, "Limits on the Transition to Gaussian Orthogonal Ensemble Behavior: Saturated Radiationless Transitions Between Strongly Coupled Potential Surfaces", *Phys. Rev. A* 40, 5935-5949 (1989).
- ++ P. Dupré, R. Jost, M. Lombardi, P.G. Green, E. Abramson, and R.W. Field, "Anomalous Behavior of the Anticrossing Density as a Function of Excitation Energy in the C_2H_2 Molecule" *J. Chem. Phys.* 152, 293-318 (1991).

- ++ K. Yamanouchi, N. Ikeda, S. Tsuchiya, D.M. Jonas, J.K. Lundberg, G.W. Adamson, and R.W. Field, "Vibrationally Highly Excited Acetylene as Studied by Dispersed Fluorescence and Stimulated Emission Pumping Spectroscopy: Vibrational Assignment of Feature States", *J. Chem. Phys.* 95, 6330-6342 (1991).
- + D.M. Jonas, Xueming Yang, and A.M. Wodtke, "Axis-Switching Transitions and the Stimulated Emission Pumping Spectrum of HCN," *J. Chem. Phys.* 97, 2284-2298 (1992).
- + D.M. Jonas, S.A.B. Solina, X. Zhao, and R.W. Field, "High Resolution VUV-Stark Measurement of the Dipole Moment of \tilde{A}^1A' HCN", *J. Chem. Phys.* 96, 7209-7217 (1992).
- + S.L. Coy, S.D. Halle, J.L. Kinsey and R.W. Field, "Pressure-Induced Rotational Energy Transfer in $H_2CO \tilde{A}^1A_2 v_4=1$: Dipolar M-Dependence with No Single-Collision Elastic Contribution", *J. Mol. Spectrosc.* 193, 340-375 (1992).
- + D.M. Jonas, S.A.B. Solina, R.W. Field, and R.J. Silbey, "Experimental Distinction of Electric and Magnetic Transition Moments," *J. Chem. Phys.* 96, 7189-7190 (1992).
- ++ J.K. Lundberg, R.W. Field, C.D. Sherrill, E.T. Seidl, Y. Xie, and H.F. Schaefer III, "Acetylene: Synergy Between Theory and Experiment," *J. Chem. Phys.* 98, 8384-8391 (1993).
- + J.K. Lundberg, Yongqin Chen, J.-P. Pique, and R.W. Field, "Ultraviolet-Optical Double Resonance Study of the Predissociated C'^1A_g State of Acetylene," *J. Mol. Spectrosc.* 156, 104-122 (1992).
- + R.L. Redington, T.E. Redington, B. Rajaram, and R.W. Field, "Excitation Spectra of 2,5-Dihydroxy-p-Benzoquinone Monomer and Hydrates," *J. Chem. Phys.* 97, 1624-1629 (1992).
- ++ D.M. Jonas, S.A.B. Solina, B. Rajaram, R.J. Silbey, R.W. Field, K. Yamanouchi, and S. Tsuchiya, "Intramolecular Vibrational Relaxation and Forbidden Transitions in the SEP Spectrum of Acetylene," *J. Chem. Phys.*, 97, 2813-2816 (1992).
- ++ D.M. Jonas, S.A.B. Solina, B. Rajaram, S.J. Cohen, R.J. Silbey, R.W. Field, K. Yamanouchi, and S. Tsuchiya, "Intramolecular Vibrational Relaxation in the SEP Spectrum of Acetylene," *J. Chem. Phys.* 99, 7350-7370 (1993).
- + J.K. Lundberg, D.M. Jonas, Yongqin Chen, Bhavani Rajaram, and R.W. Field, "Rotationally Resolved UV-UV Double Resonance Study of the Nonplanar \tilde{E}^1A_2 State of Acetylene," *J. Chem. Phys.* 97, 7180-7196 (1992).
- + P. Dupré and P.G. Green, "Characterization of a Large Singlet-Triplet Coupling Matrix Element in the Acetylene Molecule," *Chem. Phys. Lett.* 212, 555-560 (1993).
- ++ X. Zhao, G.W. Adamson, and R.W. Field, "The $HCO \tilde{B}^2A' \leftarrow \tilde{X}^2A'$ System: Fluorescence Excitation and Stimulated Emission Pumping Spectra", *J. Mol. Spectrosc.* 160, 11-38 (1993).
- + Z. J. Jakubek, N.A. Harris, R.W. Field, J.A. Gardner, and E. Murad, "The Ionization Potentials of CaF and BaF," *J. Chem. Phys.* 100, 622-627 (1994).
- +,++ Li Li and R.W. Field, "Continuous Wave Perturbation-Facilitated Optical-Optical Double Resonance Spectroscopy of Na_2 and Li_2 ," in Molecular Dynamics and Spectroscopy by Stimulated Emission Pumping pp. 251-278 (eds. H.-L. Dai and R.W. Field), World Scientific, 1995.
- + D.M. Jonas, C.A. Rogaski, A.M. Wodtke, and X. Yang, "High Resolution Spectroscopy of Chemical Isomerization: Stimulated Emission Pumping of HCN", in *Molecular Dynamics and Spectroscopy by Stimulated Emission Pumping*, p. 513-542 (eds. H.-L. Dai and R.W. Field), World Scientific, 1995.

- + S.L. Coy, D. Chasman, and R.W. Field, "Extracting Dynamical Information from Complex and Congested Spectra: Statistics, Pattern Recognition and Parismonious Trees", in Molecular Dynamics and Spectroscopy by Stimulate Emission Pumping pp. 891-942 (eds. H.-L.Dai and R.W. Field), World Scientific, 1995.
- + C. Kittrell, "Stimulated Emission Pumping by Fluorescence Dip: Experimental Methods," in *Molecular Dynamics and Spectroscopy by Stimulated Emission Pumping*, p. 109-148, (H.-L. Dai and R.W. Field, eds.), World Scientific, 1995.
- + I. Lundberg, "The SEP Spectrum of Acetylene: Symmetry Properties and Isomerization," in *Molecular Dynamics and Spectroscopy by Stimulated Emission Pumping*, p. 791-854, (H.-L. Dai and R.W. Field, eds.), World Scientific, 1995.
- ++ K. Yamanouchi, J. Miyawaki, S. Tsuchiya, D.M. Jonas, and R.W. Field, "New Scheme for Extracting Molecular Dynamics from Spectra: Case Study on Vibrationally Highly Excited Acetylene," *Laser Chemistry*, 14, 183-190 (1994).
- ++ K. Yamanouchi and R.W. Field, "Short Time Dynamics of Highly Excited Small Polyatomic Molecules Extracted from Laser Spectra," *Laser Chemistry* 16, 31-41 (1995).
- +,++ R.W. Field, S.L. Coy, and S.A.B. Solina, "Pure Sequence Vibrational Spectra of Small Polyatomic Molecules: Feature State Assignments, Sequential Dynamics, Energy and Time Scaling, and the Bag-of-Atoms Limit," *Progress of Theoretical Physics Supplement*, 116, 143-166 (1994).
- + G. M. Schmid, S. L. Coy, R. W. Field, and R.J. Silbey, "Duffing's Oscillator and the Normal to Local Mode Transition in AB₂ Triatomic Molecules", *Chem. Phys. Lett.* 219, 331-338 (1994).
- + G.M. Schmid, S.L. Coy, R.W. Field, and R.J. Silbey, "The Normal to Local Mode Transition in AB₂ Triatomic Molecules: The Susceptibility of Eigenstates to Symmetry Breaking Perturbations", *J. Chem. Phys.* 101, 869-875 (1994).
- ++ J.C. Bloch, R.W. Field, G.E. Hall, and T.J. Sears, "Time-Resolved Frequency Modulation Spectroscopy of Photochemical Transients," *J.Chem. Phys* 101, 1717-1720 (1994).
- ++ T.J. Sears, M. Wu, G.E. Hall, B.-C. Chang, G. Hansford, J.C. Bloch, and R.W. Field, "Infrared and Near Infrared Transient Absorption Spectroscopy of Molecular Free Radicals," in *Laser Techniques for State-Selected and State-to-State Chemistry*, Los Angeles, CA, Jan. 27-29, 1994, SPIE Proceedings Series No. 2124, SPIE Press, Washington (1994).
- + P. Dupré, P.G. Green, and R.W. Field, "Quantum Beat Spectroscopic Studies of Zeeman Anticrossings in the A¹A_u State of the Acetylene Molecule (C₂H₂)," *Chem. Phys.* 196, 211-238 (1995).
- + G.M. Schmid, S.L. Coy, R.W. Field, and R.J. Silbey, "How Robust are Molecular Properties? Stability Criterion for Eigenstates," *J. Chem. Phys.* 102, 337-345 (1995).
- + R.W. Field, H. Ishikawa, B. Rajaram, J. Wang, Y. Ohshima, and Y.-T. Chen, "Caught in the Act of Isomerization," 38th Welch Conference on Chemical Dynamics of Transient Species, pp. 259-284, 1994.
- ++ S.A.B. Solina, J.P. O'Brien, R.W. Field, and W.F. Polik, "Dispersed Fluorescence Spectra of Acetylene from the A¹A_u Origin: Recognition of Polyads and Test of Multiresonant Effective Hamiltonian Model for the X-¹Σ⁺ State," *J. Phys. Chem.* 100, 7797-7809 (1996).
- ++ S.A.B. Solina, J.P. O'Brien, R.W. Field, and W.F. Polik, "The Acetylene S₀ Surface: From Dispersed Fluorescence Spectra to Polyads to Dynamics", *Berichte der Bunsengesellschaft für Physikalische Chemie* 99, 555-560 (1995).

- + H. Ishikawa, Y.-T. Chen, Y. Ohshima, B. Rajaram, J. Wang, and R.W. Field, "Stimulated Emission Pumping Spectroscopy of HCP near the Isomerization Barrier: $E_{\text{VIB}} \leq 25,315 \text{ cm}^{-1}$ ", J. Chem. Phys. 105, 7383-7401 (1996).
- +,++ David J. Nesbitt and Robert W. Field, "Vibrational Energy Flow in Highly Excited Molecules: Role of Intramolecular Vibrational Redistribution," J. Phys. Chem. 100, 12,735-12,756 (1996).
- +,++ R.W. Field, J.P. O'Brien, M.P. Jacobson, S.A.B. Solina, W.F. Polik, and H. Ishikawa, "Intramolecular Dynamics in the Frequency Domain," Adv. in Chem. Phys. (Solvay Conference 1995), 1997.
- ++ M. Abbouti Tamsamani, M. Herman, S.A.B. Solina, J.P. O'Brien, and R.W. Field, "Highly Vibrational Excited $^{12}\text{C}_2\text{H}_2$ in the $\tilde{X}^1\Sigma_g^+$ State: Complementarity of Absorption and Dispersed Fluorescence Spectra," J. Chem. Phys. 105, 11357-11359 (1996).
- + S. Drucker, J.P. O'Brien, P. Patel, and R.W. Field, "The Effects of Triplet Perturbers in Photophysical Processes in the \tilde{A}^1A_u state of Acetylene," J. Chem. Phys. 100, 0000-0000 (1997).
- ++ Shih-Huang Lee, I-Chia Chen, G.W. Adamson, and R.W. Field, "The Fluorescence Excitation Spectrum of $\text{HCO } \tilde{B}^2A' - \tilde{X}^2A' (v_1^0)$ Band," J. Mol. Spectrosc. 100, 0000-0000 (1996).
- + H. Ishikawa, Chioko Nagao, Naohiko Mikami, and R.W. Field, "Observation of the 'Isomerization States' of HCP by Stimulated Emission Pumping Spectroscopy: Comparison between Theory and Experiment," J. Chem. Phys. 100, 0000-0000 (1997).
- + S.J. Humphrey, C.G. Morgan, A.M. Wodtke, K.L. Cunningham, S. Drucker, and R.W. Field, "Laser Excited Metastable States of Acetylene in the 5.5 - 5.7eV Region," J. Chem. Phys.

C. Ph.D. Theses (Since 1986) Resulting from AFOSR Support

The research of the Co-PI's is supported by AFOSR and DOE grants in the general area of the spectra of vibrationally highly excited polyatomic molecules. The distinction between these two funded projects is discussed in the original research proposals. We list below all of our Ph.D. Theses relevant to the research described in the present project; AFOSR support is denoted by (+) and DOE support by (++).

+	9/86	Vaccaro, Patrick H.	Spectroscopy and Kinetics of Highly Excited Formaldehyde
(+,++)	9/87	Cameron, Stewart M.	Photon Echo Studies of Quantum Diffractive Superposition State Scattering Kernels
++	7/88	Chen, Yongqin	Spectroscopic Studies of Highly Excited Acetylene
+	8/89	Green, Peter G.	Acetylene Near Dissociation: Novel Effects of External Fields
+	1/90	Halle, Scott D.	Spectroscopic Studies of Collision-Induced Energy Transfer and Intramolecular Dynamics of Formaldehyde
+,++	2/92	Lundberg, James K.	Double Resonance Studies of Electronically Excited Acetylene
+,++	6/92	Jonas, David M.	Spectroscopy of Vibrationally Hot Molecules: Hydrogen Cyanide and Acetylene

++	7/94	Adamson, George W.	The Spectroscopy of the Formyl Radical
(+)	9/94	Schmid, Günther M.	Dynamical Symmetry Breaking in Molecules and Molecular Aggregates.
+	1/95	Rajaram, Bhavani	Optical-Optical Double Resonance Study of the $3^1A'$ State of HCP
(+)	2/95	Gittins, Christopher	Electronic Structure and Electronic-Vibrational Energy Exchange in Rydberg States of Calcium Monofluoride
++	5/96	Solina, Stephani	Molecular Dynamics in Acetylene: From Spectra to Polyads to Dynamics
(++)	5/96	Bloch, Jonathan	Extending Frequency Modulation Spectroscopy: Sensitive and Selective High Resolution Laser Absorption in the Visible and Ultraviolet